

## Outlineoffungi.org - Note 298, [Neoacrodontiella](#)

[Neoacrodontiella](#) Crous & M.J. Wingf.

The monotypic genus [Neoacrodontiella](#) was introduced by [Crous et al. \(2019a\)](#) with *N. eucalypti* as the type species. [Neoacrodontiella](#) is somewhat reminiscent of *Acrodontiella* ([Seifert et al. 2011](#)), though distinct in that it forms sporodochia, and the conidiogenous loci are flattened and more prominent than in *Acrodontiella*, with conidia also having prominently truncate hila ([Crous et al. 2019a](#)) (M. Erdoğan).

### Reference

- Crous PW, Carnegie AJ, Wingfield MJ, Sharma R et al. 2019a – Fungal Planet description sheets, 868–950. *Persoonia* 42, 291–473. [Doi 10.3767/persoonia.2019.42.11](https://doi.org/10.3767/persoonia.2019.42.11)
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011 – The Genera of Hyphomycetes. CBS Biodiversity 9, 1–997. Westerdijk Fungal Biodiversity Institute, Utrecht, Netherlands. <https://www.nhbs.com/the-genera-of-hyphomycetes-book>