

Outlineoffungi.org - Note 481, [Sporidesmiella](#)

[Sporidesmiella](#) P.M. Kirk

[Kirk \(1982\)](#) introduced [Sporidesmiella](#) with *S. claviformis* as type species; the genus is characterized by conidia that are solitary, acrogenous, mostly distoseptate, pale olivaceous brown or subhyaline and produced by a monoblastic, terminal, integrated, indeterminate, enteroblastic percurrent elongated conidiogenous cell. This genus was accepted in *Junewangiaceae* by [Luo et al. \(2019\)](#) and [Dong et al. \(2020\)](#) based on multi-locus phylogeny and distinct morphology (W. Dong & M. Erdoğdu).

Reference

- Kirk PM. 1982 – New or interesting microfungi VI. *Sporidesmiella* gen. nov. (Hyphomycetes). Transactions of the British Mycological Society 79, 479–489.
[Doi 10.1016/S0007-1536\(82\)80040-5](#)
- Luo ZL, Hyde KD, Liu JK, Maharachchikumbura SSN et al. 2019 – Freshwater *Sordariomycetes*. Fungal Diversity 99(1), 451–660. [Doi 10.1007/s13225-019-00438-1](#)
- Dong W, Hyde KD, Doilom M, Yu XD et al. 2020 – *Pseudobactrodesmium* (*Dactylosporaceae*, *Eurotiomycetes*, *Fungi*) a Novel lignicolous genus. Frontiers in Microbiology 11 (465), 1–13.
[Doi 10.3389/fmicb.2020.00456](#)
- Dong W, Wang B, Hyde KD, McKenzie EHC et al. 2020 – Freshwater *Dothideomycetes*. Fungal Diversity 105(1), 319–575. [Doi 10.1007/s13225-020-00463-5](#)