

Outlineoffungi.org - Note 562 *Achrochaeta* - morph notes

Achrochaeta Réblová & Hern.-Restr. (H)

Achrochaeta species are saprobes on decaying wood and bark (Hughes & Kendrick 1968, Réblová et al. 2019). The monotypic genus *Achrochaeta* was introduced by Réblová et al. (2019) to accommodate *Chaetosphaeria talbotii*. *Achrochaeta* produce hyphomycetous asexual morphs which lack setae and possess cylindrical-clavate conidia (Réblová et al. 2019). No sexual morph was reported.

References

- Hughes SJ, Kendrick WB (1968) New Zealand fungi 12. *Menispora*, *Codinaea*, *Menisporopsis*. New Zealand Journal of Botany, 6:323–375. doi.org/10.1080/0028825X.1968.10428818
- Réblová M, Nekvindová J, Kolařík M, Hernández-Restrepo M (2021) Delimitation and phylogeny of *Dictyochaeta*, and introduction of *Achrochaeta* and *Tubulicolla*, genera nova. Mycologia, 113(2):390–433. doi.org/10.1080/00275514.2020.1822095

Entry by

Rekhani Hansika Perera, Center of Excellence in Fungal Research, Mae Fah Luang University, Muang, Chiang Rai, Thailand.

(Edited by **Kevin D Hyde**)