

Outlineoffungi.org - Note 574 [Zaanenomyces](#)

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#)

[Zaanenomyces](#) Crous & Osieck

[Zaanenomyces](#) was established in [Tubeufiaceae](#) ([Tubeufiales](#), [Dothideomycetes](#)) by [Crous et al. \(2021\)](#) based on morphological and phylogenetic evidence. This genus includes three species, viz. *Zaanenomyces moderatricis-academiae*, [Z. quadripartis](#) and *Z. versatilis*. All were collected from dead culms of *Juncus* spp. in the Netherlands, presumably saprobes. This hyphomycetous genus can be distinguished from other genera in [Tubeufiaceae](#) by their solitary, narrowly obclavate, straight to flexuous, septate and hyaline conidia, with subobtuse apex and obconically truncate base.

References

Crous PW, Osieck ER, Jurjevi Ž, Boers J, et al. 2021 – Fungal planet description sheets: 1284–1382. *Persoonia* 47, 178–374. <https://doi.org/10.3767/persoonia.2021.47.06>

Entry by

Yong-Zhong Lu, School of Food and Pharmaceutical Engineering, Guizhou Institute of Technology, Guiyang 550003, P.R. China.

(Edited by: **Chayanard Phukhamsakda**)

Published online 26 September 2022