

## Outlineoffungi.org - Note 84, [Bryopelta](#)

[Bryopelta](#) Döbbeler & Poelt

[Döbbeler & Poelt \(1978\)](#) introduced [Bryopelta](#) within *Pleosporaceae* to accommodate *B. variabilis*. [Bryopelta](#) is characterized by erumpent ascomata at maturity, with peridium cells of *textura angularis* to *textura porrecta*, and ascospores generally 1-septate and rarely 1–3-septate ([Döbbeler et al. 1978](#)). Subsequently, [Lumbsch & Huhndorf \(2010\)](#) referred this genus to *Dothideomycetes* genera *incertae sedis*. [Li et al. \(2014\)](#) then accommodated [Bryopelta](#) to *Mycosphaerellaceae* (M. Erdoğan).

### Reference

- Döbbeler P. 1978 – Moosbewohnende Ascomyceten I. Die pyrenocarpen, den Gametophyten besiedelnden Arten. Mitteilungen aus der Botanischen Staatssammlung München 14, 1–360. <https://www.biodiversitylibrary.org/part/194204>
- Lumbsch HT, Huhndorf SM. 2010 – Outline of *Ascomycota*-2009. Fieldiana Life and Earth Sciences 1, 1–60. [Doi 10.3158/1557.1](https://doi.org/10.3158/1557.1)
- Li WJ, Bhat JD, Hyde KD, Wang Y. 2014 – Towards a natural classification of *Dothideomycetes* 4: The genera *Bryopelta*, *Bryorella*, *Bryosphaeria*, *Lophiosphaerella* and *Maireella* (*Dothideomycetes incertae sedis*). Phytotaxa 176, 28–41. [Doi 10.11646/phytotaxa.176.1.6](https://doi.org/10.11646/phytotaxa.176.1.6)